arXiv:1212.5534v1 [math-ph] 21 Dec 2012

Perturbed GUE Minor Process and Warren's Process with Drifts

Patrik L. Ferrari* René Frings**

21. December 2012

Abstract

We consider the minor process of (Hermitian) matrix diffusions with constant diagonal drifts. At any given time, this process is determinantal and we provide an explicit expression for its correlation kernel. This is a measure on the Gelfand-Tsetlin pattern that also appears in a generalization of Warren's process [26], in which Brownian motions have level-dependent drifts. Finally, we show that this process arises in a diffusion scaling limit from an interacting particle system in the anisotropic KPZ class in $2+1$ dimensions introduced in [4].

1 Introduction

In this paper we determine a determinantal point process living on the Gelfand-Tsetlin cone GT_N,

$$\mathrm{GT}_N = \{(x^1, x^2, \ldots, x^N) \in \mathbb{R}^1 \times \mathbb{R}^2 \times \cdots \times \mathbb{R}^N : x_k^{n+1} \leq x_k^n \leq x_{k+1}^{n+1}\} \quad (1)$$

arising both from random matrices diffusions and and interacting particle systems. An element $x \in \mathrm{GT}_N$ is called a Gelfand-Tsetlin pattern. Here is a graphical representation of $x \in \mathrm{GT}_4$, which illustrates the interlacing

---

*Institute for Applied Mathematics, Bonn University, Endenicher Allee 60, 53115 Bonn, Germany. E-mail: `ferrari@uni-bonn.de`

**Institute for Applied Mathematics, Bonn University, Endenicher Allee 60, 53115 Bonn, Germany. E-mail: `frings@uni-bonn.de`

condition on x,

$$\begin{tikzcd}
x_1^4 \arrow[dr, phantom, "\leq" sloped] & & x_2^4 \arrow[dl, phantom, "\geq" sloped] \arrow[dr, phantom, "\leq" sloped] & & x_3^4 \arrow[dl, phantom, "\geq" sloped] \arrow[dr, phantom, "\leq" sloped] & & x_4^4 \arrow[dl, phantom, "\geq" sloped] \\
& x_1^3 \arrow[dr, phantom, "\leq" sloped] & & x_2^3 \arrow[dl, phantom, "\geq" sloped] \arrow[dr, phantom, "\leq" sloped] & & x_3^3 \arrow[dl, phantom, "\geq" sloped] & \\
& & x_1^2 \arrow[dr, phantom, "\leq" sloped] & & x_2^2 \arrow[dl, phantom, "\geq" sloped] & & \\
& & & x_1^1 & & &
\end{tikzcd}$$

Measures on Gelfand-Tsetlin patterns naturally appear in several fields of mathematics like (a) random matrix theory [2, 18, 14, 23] where the question of universality was recently approached in [20], (b) random tiling problems [7, 23, 22], (c) representation theory [8, 9], and (d) interacting particle systems [4, 21, 22] and diffusions [26, 27]. Probably the most famous example which belongs to more than one of these classes is the Aztec diamond. Indeed, the measure on the Aztec diamond both comes from a random tiling problem [11] and can be obtained through a Markov chain [21] on a (discrete) Gelfand-Tsetlin pattern, which itself can be seen as a special case of the more general Markov chain construction in [4]. A continuous space analogue is for instance Warren's process, a system of interacting standard Brownian diffusions [26, 27]. Recently, the link between the "drift-less" case of [4] and Warren's process has been studied in [15].

In this paper we consider GUE matrix diffusions with drifts. Its eigenvalue process at a fixed time is a determinantal point process and we explicitly determine its correlation kernel. We then show that the point process arises from a diffusion scaling limit of an interacting particle system in the anisotropic KPZ class in 2+1 dimensions [5, 4], as well as in a generalization of Warren's process [26] if we let the Brownian motions to have level-dependents drifts.

For the zero-drift case, the GUE minor process was described in [18], while Warren's process was introduced in [26], and it is known that the two processes coincide along "space-like" paths [12], they both are Markovian and determinantal, but in the whole "space-time" the two processes are different [1]. Here we focus on the fixed-time process although the connection will certainly hold along "space-like" paths as for the zero-drift case.

Matrix diffusions

The first model we study on GT_N is a variant of GUE minor process which has been introduced in [18]. Consider an $N \times N$ Hermitian matrix H with eigenvalues $\lambda_1^N \leq \cdots \leq \lambda_N^N$. Denote by H^n the submatrix obtained by keeping the first n rows and columns of H, and its ordered eigenvalues by $\lambda_1^n \leq \cdots \leq \lambda_n^n$. The collection of all these eigenvalues $(\lambda^1, \ldots, \lambda^N)$ then forms a Gelfand-Tsetlin pattern, with $\lambda^n = (\lambda^n_1, \dots, \lambda^n_n)$. In this paper we take $H(t)$ to be a GUE matrix diffusion perturbed by a deterministic drift matrix $M = \operatorname{diag}(\mu_1, \dots, \mu_N)$, i.e., we consider $G(t) = H(t) + tM$ with H evolving as standard GUE Dyson's Brownian Motion starting from 0. The eigenvalues point process ξ has support on $\mathbb{R} \times \{1, \dots, N\}$ and it is given by

$$\xi(\mathrm{d}x, m) = \sum_{1 \le k \le n \le N} \delta_{n,m} \delta_{\lambda^n_k}(\mathrm{d}x) \tag{2}$$

and their correlation functions are given as follows (for the proof, see Section 2).

Theorem 1. *For a fixed time $t > 0$ consider the eigenvalues' point process on the N submatrices of $H(t)$. Then, its m-point correlation function ϱ^m_t is given by*

$$\varrho^m_t((x_1, n_1), \dots, (x_m, n_m)) = \det[K_t((x_i, n_i), (x_j, n_j))]_{1 \le i,j \le m}, \tag{3}$$

with $(x_i, n_j) \in \mathbb{R} \times \{1, \dots, N\}$ and correlation kernel

$$K_t((x, n), (x', n')) = -\phi^{(n,n')}(x, x') + \sum_{k=1}^{n'} \Psi^{n,t}_{n-k}(x) \Phi^{n',t}_{n'-k}(x'), \tag{4}$$

where[1]

$$\phi^{(n,n')}(x, x') = \frac{(-1)^{n'-n}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z \, \frac{\mathrm{e}^{z(x'-x)}}{(z - \mu_{n+1}) \cdots (z - \mu_{n'})} \mathbb{1}_{[n<n']}, \tag{5}$$

$$\Psi^{n,t}_{n-k}(x) = \frac{(-1)^{n-k}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z \, \mathrm{e}^{tz^2/2 - xz} \frac{(z - \mu_1) \cdots (z - \mu_n)}{(z - \mu_1) \cdots (z - \mu_k)}, \tag{6}$$

$$\Phi^{n,t}_{n-\ell}(x) = \frac{(-1)^{n-\ell}}{2\pi \mathrm{i}} \oint_{\Gamma_{\mu_1, \dots, \mu_N}} \mathrm{d}w \, \mathrm{e}^{-tw^2/2 + xw} \frac{(w - \mu_1) \cdots (w - \mu_{\ell-1})}{(w - \mu_1) \cdots (w - \mu_n)} \tag{7}$$

with $\mu_- < \min\{\mu_1, \dots, \mu_N\}$.

Remark 1.1. *The integral for $\phi^{(n,n')}$ in (5) is only well-defined for $n' - n > 1$. For $n' - n = 1$ we set $\phi^{(n-1,n)}(x, x') := \phi_n(x, x') = \mathrm{e}^{\mu_n(x'-x)} \mathbb{1}_{[x>x']}$ instead.*

---

[1] For a set S, the notation $\frac{1}{2\pi \mathrm{i}} \oint_{\Gamma_S} \mathrm{d}w \, f(w)$ means that the integral is taken over any positively oriented simple contour that encloses only the poles of f belonging to S.

Warren's process with drifts

Our second model is Warren's process with drift which describes the dynamics of a system of Brownian motions $\{B_k^n, 1 \leq k \leq n \leq N\}$ on GT_N, where B_1^1 is a standard Brownian motion with drift μ_1 starting from the origin. The Brownian motions B_1^2 and B_2^2 are Brownian motions with drift μ_2 conditioned to start at the origin and, whenever they touch B_1^1, they are reflected off B_1^1. Similarly for $n \geq 2$, B_k^n is a Brownian motion with drift μ_n conditioned to start at the origin and being reflected off by B_k^{n-1} (for $k \leq n-1$) and B_{k-1}^{n-1} (for $k \geq 2$). The process with $\mu_1 = \cdots = \mu_N = 0$ was introduced and studied by Warren in [26]. In this way we obtain a system of Brownian motions with drifts that lies in GT_N. The correlation functions of this process at a fixed time agree with those of the perturbed GUE minor process (the proof is in Section 3).

Theorem 2. *For a fixed time $t > 0$ consider the point process of the positions of the Brownian motions $\{B_k^n(t) : 1 \leq k \leq n \leq N\}$ described above. Then, its m-point correlation function ϱ_t^m is also given by* (3)*.*

Interacting particle system

Finally we introduce a discrete model which gives rise to Warren's process with drifts under a diffusion scaling limit. This model is a generalization of TASEP with particle-dependent jump rates [5] to the $2+1$ dimensional particle system with Markov dynamics introduced in [4]. We denote by $x_k^n \in \mathbb{Z}$ the position of a particle labeled by (k, n), with $1 \leq k \leq n \leq N$, and call n the "level" of the particle. Particle (k, n) performs a continuous time random walk with one-sided jumps (to the right) and with rate v_n. Particles with smaller level evolve independently from the ones with higher level like in the Brownian motion model described above. More precisely, the interaction between levels is the following: (a) if particle (k, n) tries to jump to x and $x_{k-1}^{n-1} = x$, then the jump is suppressed, and (b) when particle (k, n) jumps from $x - 1$ to x, then all particles labeled by $(k + \ell, n + \ell)$ (for some $\ell \geq 1$) which were at $x - 1$ are forced to jump to x too. This is a particle system with state space in a discrete Gelfand-Tsetlin pattern.

Consider the diffusion scaling with appropriate scaled jump rates

$$t = \tau T, \quad x_k^n = \tau T - \sqrt{T}\lambda_k^n, \quad v_n = 1 - \frac{\mu_n}{\sqrt{T}}. \tag{8}$$

Then, in the $T \to \infty$ limit, the particle process $\{x_k^n(t)\}$ converges to the GUE minor process with drift $\{\lambda_k^n(\tau)\}$.

More precisely, let us denote by $\widetilde{\mathbb{P}}^v$ the probability measure on these particles with jump rates $v = (v_1, \ldots, v_N)$ given in (8). We fix $\tau > 0$ and set

$$\nu_T(A) = \widetilde{\mathbb{P}}^v\left(-\frac{x_k^n(\tau T) - \tau T}{\sqrt{T}} \in A_k^n \text{ for all } 1 \leq k \leq n \leq N\right) \tag{9}$$

where $A_k^n \subseteq \mathbb{R}$ are Borel sets, $A = \prod_{1 \leq k \leq n \leq N} A_k^n$. Moreover, we define

$$\nu(A) = \mathbb{P}^\mu\big(\lambda_k^n(\tau) \in A_k^n \text{ for all } 1 \leq k \leq n \leq N\big) \tag{10}$$

where $\mathbb{P}^\mu$ is the GUE minor measure with drift $\mathrm{diag}(\mu_1, \ldots, \mu_N)$. In Section 4 we show the following result.

Theorem 3. *As $T \to \infty$, ν_T converges to ν in total variation, i.e.,*

$$\lim_{T\to\infty} \sup_{\substack{A \subseteq \mathbb{R}^{N(N+1)/2}, \\ A \,\text{Borel}}} |\nu_T(A) - \nu(A)| = 0. \tag{11}$$

In particular, $\nu_T \to \nu$ weakly.

Acknowledgments

This work was supported by the German Research Foundation via the SFB611–A12 project.

2 GUE minor process with drift

2.1 Model and measure

Let $(H(t) : t \geq 0)$ be a process on the $N \times N$ Hermitian matrices defined by

$$H_{k\ell}(t) = \begin{cases} b_{kk}(t) + \mu_k t, & \text{if } 1 \leq k \leq N, \\ \frac{1}{\sqrt{2}}(b_{k\ell}(t) + \mathrm{i}\tilde{b}_{k\ell}(t)), & \text{if } 1 \leq k < \ell \leq N, \\ \frac{1}{\sqrt{2}}(b_{k\ell}(t) - \mathrm{i}\tilde{b}_{k\ell}(t)), & \text{if } 1 \leq \ell < k \leq N, \end{cases} \tag{12}$$

where $\{b_{kk}, b_{k\ell}, \tilde{b}_{k\ell}\}$ are independent one-dimensional standard Brownian motions[2]. Denote by $M = \mathrm{diag}(\mu_1, \ldots, \mu_N)$ the diagonal drift matrix added to the matrix H. Then, the probability measure on these matrices at time t is given by

$$\mathbb{P}(H \in \mathrm{d}H) = \mathrm{const} \times \exp\left(-\frac{\mathrm{Tr}(H - tM)^2}{2t}\right)\mathrm{d}H \tag{13}$$

---

[2] Here, standard Brownian motions start from 0 and are normalized to have variance t at time t.

where $\mathrm{d}H = \prod_{i=1}^N \mathrm{d}H_{ii} \prod_{1\leq j<k\leq N} \mathrm{d}\,\mathrm{Re}(H_{j,k})\mathrm{d}\,\mathrm{Im}(H_{j,k})$ and const is the normalization constant.

Since we are interested in the statistics of the eigenvalues' minors at time t, we first determine the measure on the eigenvalues of the $N \times N$ matrix.

Lemma 2.1. *Assume that $\mu_1, \dots, \mu_N$ are all distinct. Then under (13), the joint probability measure of the eigenvalues $\lambda_1, \dots, \lambda_N$ of H is given by*

$$\mathbb{P}(\lambda_1 \in \mathrm{d}\lambda_1, \dots \lambda_N \in \mathrm{d}\lambda_N) \\ = \mathrm{const} \times \left(\prod_{k=1}^N \mathrm{e}^{-(\lambda_k - t\mu_k)^2/(2t)}\right) \frac{\Delta(\lambda_1, \dots, \lambda_N)}{\Delta(\mu_1, \dots, \mu_N)} \mathrm{d}\lambda_1 \cdots \mathrm{d}\lambda_N \quad (14)$$

with const *a normalization constant and $\Delta(x_1, \dots, x_m) = \prod_{1\leq i<j\leq m}(x_j - x_i)$ the Vandermonde determinant.*

Remark 2.2. If $\mu_1, \dots, \mu_N$ are not all distinct, we have to take limits in (14). For instance, if $\mu_1 = \cdots = \mu_N \equiv \mu$, then

$$\mathbb{P}(\lambda_1 \in \mathrm{d}\lambda_1, \dots \lambda_N \in \mathrm{d}\lambda_N) \\ = \mathrm{const} \times \left(\prod_{k=1}^N \mathrm{e}^{-(\lambda_k - t\mu)^2/(2t)}\right) \Delta^2(\lambda_1, \dots, \lambda_N)\, \mathrm{d}\lambda_1 \cdots \mathrm{d}\lambda_N. \quad (15)$$

Proof of Lemma 2.1. We diagonalize $H = U\Lambda U^*$ with a unitary matrix U and the diagonal matrix $\Lambda = \mathrm{diag}(\lambda_1, \dots, \lambda_N)$. Then,

$$\mathrm{e}^{-\operatorname{Tr}(H-tM)^2/(2t)}\mathrm{d}H = \mathrm{const} \times \mathrm{e}^{-\operatorname{Tr}(U\Lambda U^* - tM)^2/(2t)} \Delta^2(\lambda)\, \mathrm{d}U\, \mathrm{d}\lambda, \quad (16)$$

where $\mathrm{d}U$ is the Haar measure on the unitary group $\mathcal{U}$. Moreover, since

$$\operatorname{Tr}(U\Lambda U^* - tM) = \operatorname{Tr}\Lambda^2 + t^2 \operatorname{Tr} M^2 - 2t \operatorname{Tr}(U\Lambda U^* M) \quad (17)$$

by integrating over $\mathcal{U}$ in (16) and using the Harish-Chandra-Itzykson-Zuber formula we obtain the desired expression. □

Now we focus on the minor process. For $1 \leq n \leq N$ let us denote by $H^n(t)$ the $n \times n$ principal submatrix of $H(t)$ which is obtained from $H(t)$ by keeping only the n first rows and columns. In particular, $H^1(t) = H_{11}(t)$ and $H^N(t) = H(t)$. We denote by $\lambda_1^n(t) \leq \cdots \leq \lambda_n^n(t)$ the ordered eigenvalues of $H^n(t)$. It is then a classical fact of linear algebra that at any time t, the process $(\lambda^1, \dots, \lambda^N)(t)$ lies in the Gelfand-Tsetlin cone of order N,

$$\mathrm{GT}_N = \{(x^1, \dots, x^N) \in \mathbb{R}^1 \times \cdots \times \mathbb{R}^N : x^n \preceq x^{n+1} \text{ for all } 1 \leq k \leq N-1\}, \quad (18)$$

where $x^n \preceq x^{n+1}$ means that x^n and x^{n+1} interlace, i.e.,

$$x_k^{n+1} \leq x_k^n \leq x_{k+1}^{n+1} \quad \text{for all } 1 \leq k \leq n. \tag{19}$$

The induced measure on $\{\lambda_k^n : 1 \leq k \leq n \leq N\}$ is the following.

Proposition 2.3. *Fix $t > 0$. Then, under the measure (13), the joint density of the eigenvalues of $\{H^n : 1 \leq n \leq N\}$ on GT_N is given by*

$$\text{const} \times \prod_{k=1}^{N} \mathrm{e}^{-t\mu_k^2/2} \prod_{k=1}^{N} \mathrm{e}^{-(\lambda_k^N)^2/(2t)} \Delta(\lambda^N) \prod_{\substack{1\leq n\leq N \\ 1\leq k\leq n}} \mathrm{e}^{\mu_n \lambda_k^n} \prod_{\substack{2\leq n\leq N \\ 1\leq k\leq n-1}} \mathrm{e}^{-\mu_n \lambda_k^{n-1}}, \tag{20}$$

where the normalization constant does not depend on $\mu_1, \ldots, \mu_N$.

Proof. We first derive (20) under the assumption that the $\mu_1, \ldots, \mu_N$ are all distinct. The case where some of the μ_i are equal is then recovered by taking the limit. We prove the statement inductively and follow the presentation in [13]. For $N = 1$, the density is clearly proportial to $\exp(-(\lambda_1^1 - \mu_1 t)^2/(2t))$. For $N \geq 2$, we consider an $N \times N$ matrix H^N distributed according to (13) which we write as

$$H^N - tM = \begin{pmatrix} H^{N-1} & w \\ w^* & x \end{pmatrix} - t \begin{pmatrix} M^{N-1} & 0 \\ 0 & \mu_N \end{pmatrix}, \tag{21}$$

where M^{N-1} denotes the $(N-1) \times (N-1)$ principal submatrix of M, $w \in \mathbb{C}^{N-1}$ is a Gaussian vector and $x \in \mathbb{R}$ is a Gaussian variable. Then we diagonalize H^{N-1}, i.e., we choose a unitary matrix U such that $H^{N-1} = U\Lambda U^*$ with $\Lambda = \operatorname{diag}(\lambda_1^{N-1}, \ldots, \lambda_{N-1}^{N-1})$ the diagonal matrix for the eigenvalues. Since the Gaussian distribution is invariant under unitary rotations and w is independent of H^{N-1}, we have

$$\begin{pmatrix} U^* & 0 \\ 0 & 1 \end{pmatrix} (H^N - tM) \begin{pmatrix} U & 0 \\ 0 & 1 \end{pmatrix} \overset{d}{=} \begin{pmatrix} \Lambda & w \\ w^* & x \end{pmatrix} - t \begin{pmatrix} U^* M^{N-1} U & 0 \\ 0 & \mu_N \end{pmatrix}, \tag{22}$$

where $\overset{d}{=}$ denotes equality in distribution. Applying the map $H^{N-1} \mapsto (\Lambda, U)$, the get that measure (13) on H^N is proportional to

$$\exp\left(-\frac{1}{2t} \operatorname{Tr}\left[\begin{pmatrix} \Lambda & w \\ w^* & x \end{pmatrix} - t \begin{pmatrix} U^* M^{N-1} U & 0 \\ 0 & \mu_N \end{pmatrix} \right]^2 \right) \Delta^2(\lambda^{N-1}) \\ \times \mathrm{d}U \, \mathrm{d}w \, \mathrm{d}x \, \mathrm{d}\lambda^{N-1}, \tag{23}$$

where $\mathrm{d}U$ is the Haar measure on the unitary group $\mathcal{U}_{N-1}$. We consider only the part of (23) that depends on U and integrate over $\mathcal{U}_{N-1}$ using the Harish-Chandra-Itzykson-Zuber formula

$$\int_{\mathcal{U}_{N-1}} \mathrm{d}U\, \mathrm{e}^{\mathrm{Tr}(\Lambda U^* M^{N-1} U)} = \mathrm{const} \times \frac{\det[\mathrm{e}^{\lambda_i^{N-1}\mu_j}]_{1\leq i,j\leq N-1}}{\Delta(\lambda^{N-1})\Delta(\mu_1,\ldots,\mu_{N-1})}. \tag{24}$$

After this integration the measure (23) reads

$$\mathrm{const} \times \mathbb{P}(\lambda^{N-1} \in \mathrm{d}\lambda^{N-1})\, \mathrm{e}^{x\mu_N - t\mu_N^2/2} \prod_{k=1}^{N-1} \mathrm{e}^{-|w_k|^2/t} \mathrm{d}x\, \mathrm{d}w. \tag{25}$$

We focus on the measure on w_k and represent the variables in polar coordinates, $w_k = r_k \mathrm{e}^{\mathrm{i}\varphi_k}$ with $r_k \in \mathbb{R}_+$ and $\varphi_k \in [0, 2\pi)$. Since the Jacobian of this transformation is given by $r_1 \cdots r_{N-1}$, we get

$$\prod_{k=1}^{N-1} \mathrm{e}^{-|w_k|^2/t} \mathrm{d}w_k = \prod_{k=1}^{N-1} r_k \mathrm{e}^{-r_k^2/t} \mathrm{d}r_k\, \mathrm{d}\varphi_k, \tag{26}$$

where $\mathrm{d}r_k$ and $\mathrm{d}\varphi_k$ are Lebesgue measures on $\mathbb{R}_+$ and $[0, 2\pi)$. Then we can express r_k and x in terms of the eigenvalues of H^{N-1} and H^N, see e. g. [13] for details,

$$\begin{aligned} r_k^2 &= -\frac{\prod_{j=1}^{N}(\lambda_k^{N-1} - \lambda_j^N)}{\prod_{j=1, j\neq k}^{N-1}(\lambda_k^{N-1} - \lambda_j^{N-1})} \mathbb{1}_{[\lambda^{N-1} \preceq \lambda^N]}, \\ x &= \mathrm{Tr}(H^N - H^{N-1}) = \sum_{i=1}^{N} \lambda_k^N - \sum_{k=1}^{N-1} \lambda_k^{N-1}. \end{aligned} \tag{27}$$

The Jacobian of the transformation $T : (r_1, \ldots, r_{N-1}, x) \mapsto \lambda^N$ is then given by

$$r_1 \cdots r_{N-1} |\det T'| = \frac{\Delta(\lambda^N)}{\Delta(\lambda^{N-1})} \mathbb{1}_{[\lambda^{N-1} \preceq \lambda^N]}, \tag{28}$$

and hence, given λ^{N-1}, we have

$$\begin{aligned} \mathrm{e}^{x\mu_N} \prod_{k=1}^{N-1} \mathrm{e}^{-|w_k|^2/t} \mathrm{d}x\, \mathrm{d}w = \prod_{k=1}^{N} \mathrm{e}^{-(\lambda_k^N)^2/(2t) + \mu_N \lambda_k^N} \prod_{k=1}^{N-1} \mathrm{e}^{(\lambda_k^{N-1})^2/(2t) - \mu_N \lambda_k^{N-1}} \\ \times \frac{\Delta(\lambda^N)}{\Delta(\lambda^{N-1})} \mathbb{1}_{[\lambda^{N-1} \preceq \lambda^N]}\, \mathrm{d}\lambda^N\, \mathrm{d}\varphi. \end{aligned} \tag{29}$$

Here we used that $2(r_1^2+\cdots+r_{N-1}^2)=\mathrm{Tr}(H^N)^2-\mathrm{Tr}(H^{N-1})^2$. Moreover, by the induction assumption for $N-1$ we have

$$\begin{aligned}\mathbb{P}(\lambda^{N-1}\in \mathrm{d}\lambda^{N-1}) = \mathrm{const}\times \prod_{k=1}^{N-1} \mathrm{e}^{-t\mu_k^2/2}\prod_{k=1}^{N-1}\mathrm{e}^{-(\lambda_k^{N-1})^2/(2t)}\Delta(\lambda^{N-1})\\ \times \prod_{\substack{1\leq n\leq N-1\\ 1\leq k\leq n}} \mathrm{e}^{\mu_n\lambda_k^n}\prod_{\substack{2\leq n\leq N-1\\ 1\leq k\leq n}}\mathrm{e}^{-\mu_n\lambda_k^{n-1}}\prod_{n=1}^{N-1}\mathrm{d}\lambda^n. \quad (30)\end{aligned}$$

Finally, inserting (29) and (30) into (25) and integrating out φ (which multiplies the measure by a finite constant) results in the claimed result (20). □

2.2 Correlation functions

Now we want to determine the correlation functions of the point process on the eigenvalues $\{\lambda_k^n : 1\leq k\leq n\leq N\}$. We proceed by writing the density in (20) as a product of determinants. Therefore we set $\phi_n(x,y)=\mathrm{e}^{\mu_n(y-x)}\mathbb{1}_{\{x>y\}}$ and introduce "virtual" variables $\lambda_n^{n-1}=\mathrm{virt}$ with the property that $\phi_n(\mathrm{virt},y)=\mathrm{e}^{\mu_n y}$. Then in (20) we have, up to a set measure zero,

$$\det[\phi_n(\lambda_i^{n-1},\lambda_j^n)]_{1\leq i,j\leq n}=\prod_{j=1}^{n}\mathrm{e}^{\mu_n\lambda_j^n}\prod_{j=1}^{n-1}\mathrm{e}^{-\mu_n\lambda_j^{n-1}}\mathbb{1}_{[\lambda^n\preceq\lambda^{n+1}]}. \quad (31)$$

Moreover, for $k=1,\ldots,N$ we set

$$\Psi_{N-k}^{N,t}(x)=\frac{\mathrm{e}^{-x^2/(2t)}}{\sqrt{2\pi t}}\,t^{-(N-k)/2}\,p_{N-k}\left(\frac{\mu_{k+1}t-x}{\sqrt{t}},\ldots,\frac{\mu_N t-x}{\sqrt{t}}\right), \quad (32)$$

where p_n are symmetric polynomials of degree n in n variables defined by $p_0\equiv 1$ and

$$p_n(x_1,\ldots,x_n)=\frac{(-1)^n}{\mathrm{i}\sqrt{2\pi}}\int_{\mathrm{i}\mathbb{R}}\mathrm{d}w\,\mathrm{e}^{w^2/2}(w-x_1)\cdots(w-x_n)\quad \text{for } n\geq 1. \quad (33)$$

Hence we have that

$$\prod_{k=1}^{N}\mathrm{e}^{-(\lambda_k^N)^2/(2t)}\Delta(\lambda^N)=\mathrm{const}\times\det\big[\Psi_{N-k}^{N,t}(\lambda_\ell^N)\big]_{1\leq k,\ell\leq N}, \quad (34)$$

which means that we can rewrite (20) as

$$\mathrm{const}\times\prod_{n=1}^{N}\det\big[\phi_n(\lambda_i^{n-1},\lambda_j^n)\big]_{1\leq i,j\leq n}\prod_{k=1}^{N}\mathrm{e}^{-t\mu_k^2/2}\det\big[\Psi_{N-k}^{N,t}(\lambda_\ell^N)\big]_{1\leq k,\ell\leq N}. \quad (35)$$

Note that by a change of variable $w = (tz - x)/\sqrt{t}$ we have

$$\Psi^{N,t}_{N-k}(x) = \frac{(-1)^{N-k}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}} \mathrm{d}z\, \mathrm{e}^{tz^2/2 - xz} (z - \mu_{k+1}) \cdots (z - \mu_N). \tag{36}$$

A measure of the form (35) has determinantal correlation functions and the kernel can be computed with Lemma 3.4 of [6], see the appendix.

2.3 Proof of Theorem 1

We prove the theorem first for $\mu_1 < \cdots < \mu_N$ and then use analytic continuation. Note that for $n = N$, the function $\Psi^{n,t}_{n-k}$ in (6) is the same as $\Psi^{N,t}_{N-k}$ in (36).

Lemma 2.4. *The following identities hold.*

(i) For all $n \in \{1, \ldots, N\}$, $k \in \mathbb{Z}$ and $t > 0$, we have $\phi_n * \Psi^{n,t}_{n-k} = \Psi^{n-1,t}_{n-1-k}$.

(ii) For $n < n'$, we have $\phi_{n+1} * \cdots * \phi_{n'} = \phi^{(n,n')}$ with $\phi^{(n,n')}$ given in (5).

Proof. Because of $\operatorname{Re} z < \mu_n$ we can exchange the two integrals,

$$\begin{aligned}
&(\phi_n * \Psi^{n,t}_{n-k})(x) \\
&\quad = \int_{-\infty}^{x} \mathrm{d}y\, \mathrm{e}^{\mu_n(y-x)} \frac{(-1)^{n-k}}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \mathrm{e}^{tz^2/2 - yz} \frac{(z-\mu_1)\ldots(z-\mu_n)}{(z-\mu_1)\ldots(z-\mu_k)} \\
&\quad = \frac{(-1)^{n-k}}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \mathrm{e}^{tz^2/2 - \mu_n x} \frac{(z-\mu_1)\ldots(z-\mu_n)}{(z-\mu_1)\ldots(z-\mu_k)} \int_{-\infty}^{x} \mathrm{d}y\, \mathrm{e}^{y(\mu_n - z)} \\
&\quad = \frac{(-1)^{n-1-k}}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \mathrm{e}^{tz^2/2 - xz} \frac{(z-\mu_1)\ldots(z-\mu_{n-1})}{(z-\mu_1)\ldots(z-\mu_k)} \\
&\quad = \Psi^{n-1,t}_{n-1-k}(x).
\end{aligned} \tag{37}$$

This proves the first statement. To show (ii), we first consider the case $n' - n = 2$. A simple calculation gives

$$\phi^{(n-2,n)}(x,x') = (\phi_{n-1} * \phi_n)(x,x') = -\left(\frac{\mathrm{e}^{\mu_n(x'-x)}}{\mu_n - \mu_{n-1}} + \frac{\mathrm{e}^{\mu_{n-1}(x'-x)}}{\mu_{n-1} - \mu_n} \right) \mathbb{1}_{[x > x']}, \tag{38}$$

which has the following contour integral representation,

$$\phi^{(n-2,n)}(x,x') = \frac{1}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \frac{\mathrm{e}^{z(x'-x)}}{(z-\mu_{n-1})(z-\mu_n)}. \tag{39}$$

For $n' - n > 2$, we get inductively that

$$\begin{aligned}
&(\phi_n * \phi^{(n,n')})(x,x') \\
&\quad = \frac{(-1)^{n'-n}}{2\pi \mathrm{i}} \int_{-\infty}^{x} \mathrm{d}y\, \mathrm{e}^{\mu_n(y-x)} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \frac{\mathrm{e}^{z(x'-y)}}{(z-\mu_{n+1})\cdots(z-\mu_{n'})} \\
&\quad = \frac{(-1)^{n'-n}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \frac{\mathrm{e}^{zx'-\mu_n x}}{(z-\mu_{n+1})\cdots(z-\mu_{n'})} \int_{-\infty}^{x} \mathrm{d}y\, \mathrm{e}^{y(\mu_n - z)} \qquad (40) \\
&\quad = \frac{(-1)^{n'-n+1}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z\, \frac{\mathrm{e}^{z(x'-x)}}{(z-\mu_n)(z-\mu_{n+1})\cdots(z-\mu_{n'})} \\
&\quad = \phi^{(n-1,n')}(x,x'),
\end{aligned}$$

where, as before, we could exchange the integrals because of $\operatorname{Re} z < \mu_n$. ☐

Next we consider the n-dimensional space V_n spanned by the set of functions

$$\{\phi_1 * \phi^{(1,n)}(x_1^0,\cdot), \ldots, \phi_{n-1} * \phi^{(n-1,n)}(x_{n-1}^{n-2},\cdot), \phi_n(x_n^{n-1},\cdot)\}. \qquad (41)$$

According to Lemma 3.4 of [6] we need to find a basis $\{\Phi_{n-k}^{n,t} : 1 \le k \le n\}$ of V_n that is biorthogonal to the set $\{\Psi_{n-k}^{n,t} : 1 \le k \le n\}$, i.e.,

$$\int_{\mathbb{R}} \mathrm{d}x\, \Psi_{n-k}^{n,t}(x) \Phi_{n-\ell}^{n,t}(x) = \delta_{k\ell}, \quad 1 \le k, \ell \le n. \qquad (42)$$

The form of the biorthogonal functions can be, with some experience, guessed from the form of the kernel [10].

Lemma 2.5. *We have:*

(i) V_n *is spanned by* $\{x \mapsto \mathrm{e}^{\mu_k x} : 1 \le k \le n\}$.

(ii) The functions $\{\Phi_{n-k}^{n,t} : 1 \le k \le n\}$ *are given by (7).*

Proof. For any $\varepsilon > 0$ we have

$$\begin{aligned}
&(\phi_k * \phi^{(k,n)})(x_k^{k-1}, x) \\
&\qquad = \int_{\mathbb{R}} \mathrm{d}y\, \mathrm{e}^{\mu_k y} \frac{(-1)^{n-k}}{2\pi \mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_{k+1}-\varepsilon} \mathrm{d}z\, \frac{\mathrm{e}^{z(x-y)}}{(z-\mu_{k+1})\cdots(z-\mu_n)}. \qquad (43)
\end{aligned}$$

We split the y-integral into one over $\mathbb{R}_+$ and one over $\mathbb{R}_-$. Then we can exchange the integrals over $\mathbb{R}_-$ and the imaginary axis provided that $\operatorname{Re} z < \mu_k$ and use $\int_{\mathbb{R}_-} \mathrm{d}y\, \mathrm{e}^{y(\mu_k - z)} = \frac{1}{z-\mu_k}$. In the same way we integrate over $\mathbb{R}_+$ taking

z such that $\mu_k < \operatorname{Re} z < \mu_{k+1}$. This gives $\int_{\mathbb{R}_+} \mathrm{d}y\, \mathrm{e}^{y(\mu_k - z)} = -\frac{1}{z-\mu_k}$. Putting these two integrals together we get

$$(\phi_k * \phi^{(k,n)})(x_k^{k-1+1}, x) = \frac{(-1)^{n-k}}{2\pi\mathrm{i}} \oint_{\Gamma_{\mu_k}} \mathrm{d}z\, \frac{\mathrm{e}^{xz}}{(z-\mu_k)(z-\mu_{k+1})\cdots(z-\mu_n)}, \tag{44}$$

which is a constant multiple of $\mathrm{e}^{\mu_k x}$. This proves (i). For (ii) we proceed similarly. Using that $1 \leq k, \ell \leq n$ we have

$$\begin{aligned}&\int_{\mathbb{R}} \mathrm{d}x\, \Psi_{n-k}^{n,t}(x) \Phi_{n-\ell}^{n,t}(x) \\ &\quad = \frac{(-1)^{k+\ell}}{(2\pi\mathrm{i})^2} \int_{\mathbb{R}} \mathrm{d}x \int_{\mathrm{i}\mathbb{R}} \mathrm{d}z \oint_{\Gamma_{\mu_\ell,\dots,\mu_n}} \mathrm{d}w\, \frac{\mathrm{e}^{tz^2 - xz}}{\mathrm{e}^{tw^2 - xw}} \frac{(z-\mu_{k+1})\cdots(z-\mu_n)}{(w-\mu_\ell)\cdots(w-\mu_n)}.\end{aligned} \tag{45}$$

When we integrate x over $\mathbb{R}_-$, we take the z-integral such that $\operatorname{Re} z < \operatorname{Re} w$, and when we integrate x over $\mathbb{R}_+$, we choose $\operatorname{Re} z > \operatorname{Re} w$. Thus, (45) reduces to

$$\frac{(-1)^{k+\ell}}{2\pi\mathrm{i}} \oint_{\Gamma_{\mu_\ell,\dots,\mu_n}} \mathrm{d}w\, \frac{(w-\mu_{k+1})\cdots(w-\mu_n)}{(w-\mu_\ell)\cdots(w-\mu_n)} = \delta_{k\ell}. \tag{46}$$

Finally, note that

$$\Phi_{n-\ell}^{n,t}(x) = \sum_{i=\ell}^{n} b_i \mathrm{e}^{\mu_i x} \quad \text{with} \quad b_i = \prod_{\substack{j=\ell \\ j\neq i}}^{n} \frac{\mathrm{e}^{-t\mu_i^2/2}}{\mu_i - \mu_j}. \tag{47}$$

which shows that the set $\{\Phi_{n-k}^{n,t} : 1 \leq k \leq n\}$ spans V_n. □

Next we verify Assumption (A) from Lemma 3.4 in [6]. Indeed,

$$\Phi_0^{n,t}(x) = \frac{1}{2\pi\mathrm{i}} \oint_{\Gamma_{\mu_n}} \mathrm{d}w\, \frac{\mathrm{e}^{-tw^2/2 + xw}}{w - \mu_n} = c_n \phi_n(x_n^{n-1}, x) \tag{48}$$

with $c_n = \mathrm{e}^{-t\mu_n^2/2} \neq 0$ for $n = 1, \dots, N$.

Finally, we can also determine the value of the normalization constant in (35), since it is given by $1/\det[M_{k\ell}]_{1\leq k,\ell\leq N}$ with

$$M_{k\ell} = (\phi_k * \cdots * \phi_N * \Psi_{N-\ell}^{N,t})(x_k^{k-1}). \tag{49}$$

Lemma 2.6. *We have* $\det M = \prod_{n=1}^{N} \mathrm{e}^{t\mu_n^2/2}$, *in particular* $\det M > 0$.

Proof. By Lemma 2.4 (i) we may write $M_{k\ell} = (\phi_k * \Psi^{k,t}_{k-\ell})(x_k^{k-1})$. Thus, for $k \geq \ell$,

$$M_{k\ell} = \int_{\mathbb{R}} \mathrm{d}y\, \mathrm{e}^{\mu_k y} \frac{(-1)^{k-\ell}}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}} \mathrm{d}z\, \mathrm{e}^{tz^2/2-yz} \, (z-\mu_{\ell+1})\cdots(z-\mu_k). \tag{50}$$

Once again, we let run the y-integral over $\mathbb{R}_-$ and $\mathbb{R}_+$ separately. In the first case we take the z-integral such that $\operatorname{Re} z < \mu_k$, in the second case such that $\operatorname{Re} z > \mu_k$. This allows us to exchange the integrals, which gives

$$M_{k\ell} = \frac{(-1)^{k-\ell}}{2\pi\mathrm{i}} \oint_{\Gamma_{\mu_k}} \mathrm{d}z\, \mathrm{e}^{tz^2/2} \, \frac{(z-\mu_{\ell+1})\cdots(z-\mu_k)}{z-\mu_k}. \tag{51}$$

Since the integrand has no poles for $k > \ell$, we have $M_{k\ell} = 0$ in this case, while for $k = \ell$ we get $M_{kk} = \mathrm{e}^{-t\mu_k^2/2}$. Thus, M is upper triangular and the claim follows. ☐

With the results of Lemma 2.4 and Lemma 2.5, Theorem 1 follows directly from Lemma 3.4 of [6].

We have shown that Theorem 1 holds when we impose $\mu_1 < \cdots < \mu_N$. In particular, the joint density (20) is given by a $N(N+1)/2$ correlation function: Let $m = N(N+1)/2$, then

$$(20) = m!\, \rho^{(m)}(\{(\lambda_k^n, n), 1 \leq k \leq n \leq N\}). \tag{52}$$

Let $M > 0$ be any fixed real number. The density (20) is analytic in each of the μ_j in $[-M, M]$, $j = 1, \ldots, N$. The same holds for the correlation kernel (take e.g., $\mu_- = -M-1$). From this it follows that also the r.h.s. of (52) is analytic in each of the variables $\mu_1, \ldots, \mu_N$. Since this holds for any M, by analytic continuation it follows that Theorem 1 holds for any given drift vector $(\mu_1, \ldots, \mu_N) \in \mathbb{R}^N$.

3 Warren's process with drifts

We have seen in Section 2 that the eigenvalues' density can be written as a product of determinants, and, in Lemma 2.6, we calculated the normalization constant, so that the probability measure on the eigenvalues reads

$$\begin{aligned}\mathbb{P}\Big(\bigcap_{1\leq k\leq n\leq N} \{\lambda_k^n \in \mathrm{d}\lambda_k^n\}\Big) &= \det\big[\Psi^{N,t}_{N-k}(\lambda_\ell^N)\big]_{1\leq k,\ell\leq N} \prod_{n=1}^N \mathrm{e}^{-t\mu_n^2/2} \\ &\times \prod_{n=1}^N \det\big[\phi_n(\lambda_i^{n-1}, \lambda_j^n)\big]_{1\leq i,j\leq n} \prod_{1\leq k\leq n\leq N} \mathrm{d}\lambda_k^n. \end{aligned} \tag{53}$$



Figure 1: The two reflection types in our system. They correspond to the boundary condition (54).

In this section we show that this measure also describes the transition probabilities for a system of Brownian motions in GT_N, starting from 0 at time $t = 0$ and arriving at $x \in \mathrm{GT}_N$ at time $t > 0$. Let us consider $\{B_k^n, 1 \leq k \leq n \leq N\}$ Brownian motions in GT_N starting from 0, with drift μ_n, and interacting as follows:

- the evolution of B_k^n does not depends on the Brownian motions with higher upper index (B_ℓ^m for $m \geq n+1$, and any ℓ),
- B_k^n is reflected off by B_k^{n-1} and B_{k-1}^{n-1}.

These reflections are sometimes called oblique reflections [25], since in the (x_k^{n-1}, x_k^n) plane (resp. (x_{k-1}^{n-1}, x_k^n) plane) the reflection direction is not the normal one, see Figure 1. Remark that the projection on $\{B_1^n, 1 \leq n \leq N\}$ differs from the process studied in [24], where the reflections are in the normal direction. If we denote by p_t the probability density in GT_N of the system of reflected Brownian motions, this kind of reflections are described by the boundary condition

$$\frac{\partial}{\partial x_k^n} p_t(x) + (\mu_{n+1} - \mu_n) p_t(x) = 0, \tag{54}$$

whenever $x_k^n = x_k^{n+1}$ or $x_k^n = x_{k+1}^{n+1}$, for $1 \leq k \leq N-1$. This is a special case of the reflection in a given direction studied in [16]. For a Brownian motion with drift μ reflected at the boundary in the direction v, the boundary conditions are the following. Denote by n the normal vector of the boundary, let v be normalized such that $n \cdot v = 1$ and let $q = v - n$. Moreover, set $D^* = n \cdot \nabla - q \cdot (\nabla - n(n \cdot \nabla))$. Then, the boundary condition are written as

$$D^* p_t = 2(\mu \cdot n) p_t, \quad \text{at the boundary.} \tag{55}$$

Specializing to our case, we get (54). This process, without drifts, was introduced by Warren in [26]. In that paper, he determined the transition probability for any initial condition and also showed that the process is well-defined when starting from 0. We here consider a system of Brownian motions with constant (bounded) drifts, which can be expressed as follows,

$$\begin{aligned}
&B_1^1(t) = \mu_1 t + b_1^1(t),\\
&B_1^n(t) = \mu_n t + b_1^n(t) - L_{B_1^{n-1}-B_1^n}(t), \quad n = 2,\dots,N,\\
&B_k^n(t) = \mu_n t + b_k^n(t) - L_{B_k^{n-1}-B_k^n}(t) + L_{B_k^n-B_{k-1}^{n-1}}(t), \quad 2 \le k < n \le N,\\
&B_n^n(t) = \mu_n t + b_n^n(t) + L_{B_n^n-B_{n-1}^{n-1}}(t), \quad n = 2,\dots,N,
\end{aligned} \tag{56}$$

where the b_k^n, $1 \le k \le n \le N$, are independent standard Brownian motions and $L_{X-Y}(t)$ is the local time accumulated at the origin by $X(t) - Y(t)$. The question of well-definedness was related with the, a priori possible, presence of triple collisions. Bounded drifts do not influence this property as can be seen by applying Girsanov's theorem like in the works [17, 19]. In this paper we do not determine the full transition probability, but rather the transition probability for the our specific initial condition.

Proposition 3.1. *Denote by* $p_t : \mathrm{GT}_N \to [0,1]$ *the transition probability density of the* $N(N+1)/2$ *Brownian motions with drift, starting from* 0 *at time* $t = 0$ *and arriving at* $x \in \mathrm{GT}_N$ *at time* $t > 0$. *Inside* GT_N, *this density satisfies the Fokker-Planck equation*

$$\frac{\partial}{\partial t} p_t(x) = \sum_{n=1}^{N} \sum_{k=1}^{n} \left(\frac{1}{2} \frac{\partial^2}{\partial (x_k^n)^2} - \mu_n \frac{\partial}{\partial x_k^n} \right) p_t(x). \tag{57}$$

Moreover, as $t \searrow 0$, *the measure* $p_t(x)\mathrm{d}x$ *becomes the Dirac measure at* 0,

$$\lim_{t \searrow 0} p_t(x)\mathrm{d}x = \prod_{1 \le k \le n \le N} \delta_{x_k^n}, \tag{58}$$

and the reflections are translated by the boundary condition

$$\frac{\partial}{\partial x_k^n} p_t(x) + (\mu_{n+1} - \mu_n) p_t(x) = 0 \tag{59}$$

for $1 \le k \le N-1$, *whenever* $x_k^n = x_k^{n+1}$ *or* $x_k^n = x_{k+1}^{n+1}$.

Proof. First observe that by setting $\tilde{\Psi}_{N-k}^{N,t}(x) = \mathrm{e}^{\mu_N x} \Psi_{N-k}^{N,t}(x)$, we can rewrite (53) as a probability measure on GT_N with density

$$p_t(x) = \det \left[\tilde{\Psi}_{N-k}^{N,t}(x_\ell^N) \right]_{1 \le k, \ell \le N} \prod_{k=1}^{N} \mathrm{e}^{-t\mu_k^2/2} \prod_{n=1}^{N-1} \prod_{k=1}^{n} \mathrm{e}^{(\mu_n - \mu_{n+1}) x_k^n}. \tag{60}$$

for $x = (x^n_k)_{1\le k\le n\le N} \in \mathrm{GT}_N$. The double product only depends on $(x^n_k)_{1\le k\le n\le N-1}$, while the determinant is a function of $(x^N_k)_{1\le k\le N}$. We have

$$\frac{1}{2}\frac{\partial^2}{\partial x^2}\,\tilde{\Psi}^{N,t}_{N-k}(x) = \frac{\partial}{\partial t}\,\tilde{\Psi}^{N,t}_{N-k}(x) + \mu_N\frac{\partial}{\partial x}\tilde{\Psi}^{N,t}_{N-k}(x) - \frac{\mu_N^2}{2}\tilde{\Psi}^{N,t}_{N-k}(x), \tag{61}$$

from which follows that

$$\frac{1}{2}\sum_{\ell=1}^{N}\frac{\partial^2}{\partial(x^N_\ell)^2}\,p_t(x) = \frac{\partial}{\partial t}\,p_t(x)+\mu_N\sum_{\ell=1}^{N}\frac{\partial}{\partial x^N_\ell}\,p_t(x)+\frac{1}{2}\left(\sum_{n=1}^{N}\mu_n^2 - N\mu_N^2\right)p_t(x). \tag{62}$$

For $k = 1,\dots,N-1$, we have

$$\frac{\partial}{\partial x^n_k}\,p_t(x) = (\mu_n-\mu_{n+1})p_t(x), \qquad \frac{\partial^2}{\partial(x^n_k)^2}\,p_t(x) = (\mu_n-\mu_{n+1})^2 p_t(x), \tag{63}$$

and thus, putting (62) and (63) together,

$$\begin{aligned}\frac{1}{2}\sum_{n=1}^{N}\sum_{k=1}^{n}\frac{\partial^2}{\partial(x^n_k)^2}\,p_t(x) &= \frac{\partial}{\partial t}\,p_t(x) + \mu_N\sum_{k=1}^{N}\frac{\partial}{\partial x^N_k}\,p_t(x)\\ &\quad + \frac{1}{2}\left(\sum_{n=1}^{N}\mu_n^2 - N\mu_N^2 + \sum_{n=1}^{N-1}n(\mu_n-\mu_{n+1})^2\right)p_t(x).\end{aligned} \tag{64}$$

Using that

$$N\mu_N^2-\sum_{n=1}^{N}\mu_n^2 = \sum_{n=1}^{N-1}n(\mu_{n+1}^2-\mu_n^2) = \sum_{n=1}^{N-1}n(\mu_n-\mu_{n+1})^2-2\sum_{k=1}^{N-1}n\mu_n(\mu_n-\mu_{n+1}) \tag{65}$$

the expression between the brackets in (64) simplifies to $2\sum n\mu_n(\mu_n-\mu_{n+1})$. On the other hand,

$$\sum_{n=1}^{N}\sum_{k=1}^{n}\mu_n\frac{\partial}{\partial x^n_k}\,p_t(x) = \sum_{n=1}^{N-1}n\mu_n(\mu_n-\mu_{n+1})p_t(x) + \mu_N\sum_{k=1}^{N}\frac{\partial}{\partial x^N_k}\,p_t(x). \tag{66}$$

Then, (57) follows from (64), (65) and (66). The initial condition (58) is satified because as $t\searrow 0$, we obtain the Dirac measure at $x^N_k = 0$ for $1\le k\le N$ and since we consider p_t on GT_N, this immediately implies that $x^k_n = 0$ for all $1\le k\le n\le N-1$. Finally, the boundary condition (59) holds trivially by (63). $\square$


$n=5$
$n=4$
$n=3$
$n=2$
$n=1$
x_1^5 x_2^5 x_3^5 x_4^5 x_5^5
x_1^4 x_2^4 x_3^4 x_4^4
x_1^3 x_2^3 x_3^3
x_1^2 x_2^2
x_1^1

Figure 2: (Left) Initial particles configuration. (Right) A possible particles configuration after some time; in this configuration, if particle $(1,3)$ tries to jump, the move is suppressed (blocked by particle $(1,2)$), while if particle $(2,2)$ jumps, then also particles $(3,3)$ and $(4,4)$ move by one unit to the right. Particles at level n have a jump rate v_n.

4 $2+1$ dynamics with different jump rates

In this section we show that the correlation functions (3) that we obtained for the GUE matrix diffusion with drifts (and for Warren's process with drifts) can be obtained as a limit from an GT_N-extension of TASEP with particle-dependent jump rates. This latter process was introduced in [4]. Before we come to the convergence result, let us describe the model.

At a fixed time t, let us denote by $x(t)=(x_k^n(t))_{1\le k\le n\le N}\in\mathrm{GT}_N$ the positions of the $N(N+1)/2$ particles at time t. We choose initial conditions $x_k^n(0)=k-n-1$ and let the particles evolve as follows: Each particle x_k^n has an independent exponential clock of rate $v_n>0$, i.e., particles on the same level have the same jump rates. When the x_k^n-clock rings, the particle jumps to the right by one, provided that $x_k^n<x_k^{n-1}-1$, otherwise we say that x_k^n is blocked by x_k^{n-1}. If the x_k^n-particle can jump, we take the largest $c\ge 1$ such that $x_k^n=x_{k+1}^{n+1}=\cdots=x_{k+c-1}^{n+c-1}$, and all c particles in this string jump to the right by one, see Figure 2 for an example). This ensures that at any time t, all the particles are in GT_N. More precisely, these dynamics imply that the particles stay in a discrete version of GT_N, namely

$$\{(x^1,x^2,\dots,x^N)\in\mathbb{Z}^1\times\mathbb{Z}^2\times\cdots\times\mathbb{Z}^N : x_k^{n+1}<x_k^n\le x_{k+1}^{n+1}\}. \tag{67}$$

The joint distribution of the particles has been calculated in Theorem 4.1 of [5]. It is given by

$$\mathrm{const}\times\det\big[\widetilde{\Psi}^{N,t}_{N-k}(x_\ell^N)\big]_{1\le k,\ell\le N}\prod_{n=1}^{N}\det\big[\widetilde{\phi}_n(x_i^{n-1},x_j^n)\big]_{1\le i,j\le n}, \tag{68}$$

where

$$\begin{aligned}\widetilde{\Psi}^{N,t}_{N-k}(x) &= \frac{1}{2\pi\mathrm{i}}\oint_{\Gamma_0} \mathrm{d}z\, \mathrm{e}^{t/z} z^{x+N-1}(1-v_{k+1}z)\cdots(1-v_N z),\\ \widetilde{\phi}_n(x,y) &= (v_n)^{y-x}\mathbb{1}_{[y\geq x]} \quad \text{and} \quad \widetilde{\phi}_n(x_n^{n-1},y) = (v_n)^y.\end{aligned} \tag{69}$$

Actually Theorem 4.1 of [5] is a statement about the marginal of a (possibly signed) measure. However, this model is the continuous time limit of a generic Markov chain introduced in Section 2 of [4], from which it follows that the measure with fully packed initial conditions $y_n = x_1^n(0) = -n$ for $1 \leq n \leq N$ is actually a probability distribution. The form of (68) follows then from the theorem by taking $a(t) = t$ and $b(t) = 0$ for all $t \geq 0$. Also note that we put the transition from time $t = 0$ to time t (which is encoded by $\mathcal{T}_{t,0}$ in the theorem) into Ψ^N_{N-k}. The correlation functions of this point process are determinantal as shown in [5]. What remains to do is the biorthogonalization for the generic jump rates.

Proposition 4.1. *Consider a system of particles on* GT_N *with fully packed initial conditions and dynamics described above. Then, at fixed time* t*, the corresponding point process has* m*-point correlation function* $\widetilde{\varrho}_t^m$ *given by*

$$\widetilde{\varrho}_t^m((x_1,n_1),\ldots,(x_m,n_m)) = \det[\widetilde{K}^v_t((x_i,n_i),(x_j,n_j))]_{1\leq i,j\leq m} \tag{70}$$

with $(x_i,n_i) \in \mathbb{R}\times\{1,\ldots,N\}$ *and correlation kernel*

$$\widetilde{K}^v_t((x,n),(x',n')) = -\widetilde{\phi}^{(n,n')}(x,x') + \sum_{k=1}^{n'} \widetilde{\Psi}^{n,t}_{n-k}(x)\widetilde{\Phi}^{n',t}_{n'-k}(x'), \tag{71}$$

where

$$\widetilde{\phi}^{(n,n')}(x,x') = \frac{1}{2\pi\mathrm{i}}\oint_{\Gamma_{0,v}} \mathrm{d}z\, \frac{1}{z^{x-x'+1}}\, \frac{z^{n'-n}}{(z-v_{n+1})\cdots(z-v_{n'})}\,\mathbb{1}_{[n<n']}, \tag{72}$$

$$\widetilde{\Psi}^{n,t}_{n-k}(x) = \frac{1}{2\pi\mathrm{i}}\oint_{\Gamma_{0,v}} \mathrm{d}z\, \frac{\mathrm{e}^{tz}}{z^{x+n+1}}\, \frac{(z-v_1)\cdots(z-v_n)}{(z-v_1)\cdots(z-v_k)}, \tag{73}$$

$$\widetilde{\Phi}^{n,t}_{n-\ell}(x) = \frac{1}{2\pi\mathrm{i}}\oint_{\Gamma_v} \mathrm{d}w\, \frac{w^{x+n}}{\mathrm{e}^{tw}}\, \frac{(w-v_1)\cdots(w-v_{\ell-1})}{(w-v_1)\cdots(w-v_n)}. \tag{74}$$

Proof. By Proposition 3.1 of [5], we have

$$\widetilde{\Psi}^{n,t}_{n-k}(x) = \frac{1}{2\pi\mathrm{i}}\oint_{\Gamma_0} \mathrm{d}w\, w^{x+k-1}\mathrm{e}^{t/w}\, \frac{(1-v_1w)\cdots(1-v_nw)}{(1-v_1w)\cdots(1-v_kw)} \tag{75}$$

for $k \geq 1$. A change of variable $z = 1/w$ then yields (73). Next we need to verify that $\{\widetilde{\Psi}^{n,t}_{n-k} : 1 \leq k \leq n\}$ is biorthogonal to $\{\widetilde{\Phi}^{n,t}_{n-\ell} : 1 \leq \ell \leq n\}$ (see Eq. (3.5) of [5]). We split the sum over $\mathbb{Z}$ into two parts, one over $x \geq 0$ and one over $x < 0$. Then,

$$\begin{aligned}\sum_{x\geq 0} & \widetilde{\Psi}^{n,t}_{n-k}(x)\widetilde{\Phi}^{n,t}_{n-k}(x) \\ &= \sum_{x\geq 0} \frac{1}{(2\pi \mathrm{i})^2} \oint_{\Gamma_v} \mathrm{d}w \oint_{\Gamma_0} \mathrm{d}z \, \frac{\mathrm{e}^{tz}}{\mathrm{e}^{tw}} \, \frac{w^{x+n}}{z^{x+n+1}} \, \frac{(z-v_{k+1})\cdots(z-v_n)}{(w-v_\ell)\cdots(w-v_n)}. \qquad (76)\end{aligned}$$

We choose Γ_0 and Γ_v such that $|w| \leq |z|$ which allows us to put the sum inside the integrals. This gives

$$\begin{aligned}\sum_{x\geq 0} & \widetilde{\Psi}^{n,t}_{n-k}(x)\widetilde{\Phi}^{n,t}_{n-k}(x) \\ &= \frac{1}{(2\pi \mathrm{i})^2} \oint_{\Gamma_v} \mathrm{d}w \oint_{\Gamma_{0,w}} \mathrm{d}z \, \frac{\mathrm{e}^{tz}}{\mathrm{e}^{tw}} \, \frac{w^{n}}{z^{n}} \, \frac{(z-v_{k+1})\cdots(z-v_n)}{(w-v_k)\cdots(w-v_\ell)} \, \frac{1}{z-w}. \qquad (77)\end{aligned}$$

For $x < 0$ we choose Γ_0 and Γ_v such that they satisfy $|w| > |z|$ which gives

$$\begin{aligned}\sum_{x< 0} & \widetilde{\Psi}^{n,t}_{n-k}(x)\widetilde{\Phi}^{n,t}_{n-k}(x) \\ &= -\frac{1}{(2\pi \mathrm{i})^2} \oint_{\Gamma_0} \mathrm{d}z \oint_{\Gamma_{v,z}} \mathrm{d}w \, \frac{\mathrm{e}^{tz}}{\mathrm{e}^{tw}} \, \frac{w^{n}}{z^{n}} \, \frac{(z-v_{k+1})\cdots(z-v_n)}{(w-v_k)\cdots(w-v_\ell)} \, \frac{1}{z-w}. \qquad (78)\end{aligned}$$

Thus,

$$\sum_{x\in\mathbb{Z}} \widetilde{\Psi}^{n,t}_{n-k}(x)\widetilde{\Phi}^{n,t}_{n-k}(x) = \frac{1}{2\pi \mathrm{i}} \oint_{\Gamma_v} \mathrm{d}w \, \frac{(w-v_{k+1})\cdots(w-v_n)}{(w-v_\ell)\cdots(w-v_n)} = \delta_{k\ell}. \qquad (79)$$

Finally, we show that $\{\widetilde{\Phi}^{n,t}_{n-\ell} : 1 \leq \ell \leq n\}$ spans the space of functions V_n. We denote by $u_1 < \cdots < u_\nu$ the different values of $v_1, \ldots, v_n$ and α_k the multiplicity of u_k, i.e., $\alpha_1 + \cdots + \alpha_\nu = n$. Then, we may write

$$\begin{aligned}\widetilde{\Phi}^{n,t}_{n-1}(x) &= \frac{1}{2\pi \mathrm{i}} \oint_{\Gamma_v} \mathrm{d}w \, \frac{w^{x+n}}{\mathrm{e}^{tw}} \, \frac{1}{(w-u_1)^{\alpha_1}\cdots(w-u_\nu)^{\alpha_\nu}} \\ &= \sum_{i=1}^{\nu} \frac{1}{(\alpha_i - 1)!} \, \frac{\mathrm{d}^{\alpha_i - 1}}{\mathrm{d}w^{\alpha_i - 1}}\bigg|_{w=u_i} \left(\frac{w^{x+n}}{\mathrm{e}^{tw}} \prod_{j\neq i} \frac{1}{(w-u_j)^{\alpha_j}} \right) \qquad (80) \\ &= \sum_{i=1}^{\nu} (u_i)^x \sum_{j=1}^{\alpha_i} c_{i,j} x^{j-1}.\end{aligned}$$

For $\ell = 2, \dots, n$, we can represent $\widetilde{\Phi}^{n,t}_{n-\ell}$ in the same way, but with exponents $\alpha_{\ell,i} \leq \alpha_i$, $1 \leq i \leq \nu$. Since $(\alpha_{k,1}, \dots, \alpha_{k,\nu}) \neq (\alpha_{\ell,1}, \dots, \alpha_{\ell,\nu})$ for $k \neq \ell$, this shows that

$$\operatorname{span}\{\widetilde{\Phi}^{n,t}_{n-\ell} : 1 \leq \ell \leq n\} = \operatorname{span}\{x \mapsto (u_i)^x x^{j-1} : 1 \leq u \leq \nu, 1 \leq j \leq \alpha_i\}, \tag{81}$$

which is V_n. ☐

We continue by establishing the convergence result under the scaling (8). Correspondingly, we rescale (and conjugate) the kernel $\widetilde{K}_t$ and define the rescaled kernel as

$$K^{\mu}_{\tau,T,\mathrm{resc}}((\xi,n),(\xi',n')) = \frac{T^{n'/2}}{T^{n/2}} \sqrt{T}\, \widetilde{K}^{\mu_T}_{\tau T}\big(([\tau T - \xi\sqrt{T}], n), ([\tau T - \xi'\sqrt{T}], n')\big) \tag{82}$$

where $[\,\cdot\,]$ denotes the integer part, and the drift v is now $\mu_T = 1 - \mu/\sqrt{T}$. Of course, T is assumed to be so large that $\mu_T > 0$ is satisfied.

Proposition 4.2. *For any fixed $L > 0$, the rescaled kernel $K^{\mu}_{\tau,T,\mathrm{resc}}$ converges, uniformly for $\xi, \xi' \in [-L, L]$, as*

$$\lim_{T\to\infty} K^{\mu}_{\tau,T,\mathrm{resc}}((\xi,n),(\xi',n')) = K^{\mu}_{\tau}((\xi,n),(\xi',n')) \tag{83}$$

with $K^{\mu}_{\tau} \equiv K_\tau$ given in (4).

Proof. Let us define the rescaled functions

$$\begin{aligned}
\phi^{(n,n')}_{T,\mathrm{resc}}(\xi,\xi') &= T^{-(n'-n+1)/2}\, \widetilde{\phi}^{(n,n')}(\tau T + \xi\sqrt{T}, \tau T + \xi'\sqrt{T}),\\
\Psi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi) &= T^{(n-k+1)/2} \mathrm{e}^{-\tau T}\, \widetilde{\Psi}^{n,\tau T}_{n-k}(\tau T + \xi\sqrt{T}),\\
\Phi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi') &= T^{-(n-k)/2} \mathrm{e}^{\tau T}\, \widetilde{\Phi}^{n,\tau T}_{n-k}(\tau T + \xi'\sqrt{T}),
\end{aligned} \tag{84}$$

where we also rescale the jump rates as in (8). We have to show that these functions converge to their analogues from (5)–(7). We first verify that $\phi^{(n,n')}_{T,\mathrm{resc}}(\xi,\xi') \to \phi^{(n,n')}(\xi,\xi')$ with $n < n'$. For $y \geq y'$, the integrand of $\widetilde{\phi}^{(n,n')}(y,y')$ in (72) has residue 0 at infinity and thus the whole integral vanishes, while for $y < y'$, there is no pole at $z = 0$ and therefore

$$\widetilde{\phi}^{(n,n')}(y,y') = \sum_{i=n+1}^{n'} v_i^{(y'-y)+(n'-n)-1} \prod_{j\neq i} \frac{1}{v_i - v_j}\, \mathbb{1}_{[y<y']}. \tag{85}$$

Hence, for its rescaled version,

$$\phi^{(n,n')}_{T,\mathrm{resc}}(\xi,\xi') = \sum_{i=n+1}^{n'} \left(1 - \frac{\mu_i}{\sqrt{T}}\right)^{(\xi-\xi')\sqrt{T}+(n'-n)-1} \prod_{j\neq i} \frac{1}{\mu_j - \mu_i}\, \mathbb{1}_{[\xi>\xi']}, \tag{86}$$

which, as $T \to \infty$, converges to

$$\sum_{i=n+1}^{n'} \mathrm{e}^{\mu_i(\xi'-\xi)} \prod_{j\neq i} \frac{1}{\mu_j - \mu_i} \mathbb{1}_{\{\xi > \xi'\}} = \frac{(-1)^{n'-n}}{2\pi\mathrm{i}} \int_{\mathrm{i}\mathbb{R}+\mu_-} \mathrm{d}z \, \frac{\mathrm{e}^{z(\xi'-\xi)}}{(z-\mu_{n+1})\cdots(z-\mu_{n'})}. \tag{87}$$

Next we show that $\Psi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi) \to \Psi^{n,\tau}_{n-k}(\xi)$ uniformly for $\xi \in [-L, L]$. We have

$$\begin{aligned} \Psi^{n,\tau,T}_{n-k,\mathrm{resc}}(\xi) &= \frac{\sqrt{T}}{2\pi\mathrm{i}} \oint_{\Gamma_{0,v}} \mathrm{d}z \, \frac{\mathrm{e}^{\tau T(z-1)}}{z^{\tau T - \xi\sqrt{T} + n + 1}} \, g(z) \\ &= \frac{\sqrt{T}}{2\pi\mathrm{i}} \oint_{\Gamma_{0,v}} \mathrm{d}z \, \mathrm{e}^{\tau T f_0(z) + \sqrt{T} f_1(z) + f_2(z)} g(z) \end{aligned} \tag{88}$$

with $f_0(z) = z - 1 - \ln z$, $f_1(z) = \xi \ln z$, $f_2(z) = -(n+1)\ln z$, and

$$g(z) = \frac{(\sqrt{T}(z-1)+\mu_1)\cdots(\sqrt{T}(z-1)+\mu_n)}{(\sqrt{T}(z-1)+\mu_1)\cdots(\sqrt{T}(z-1)+\mu_k)}. \tag{89}$$

A Taylor expansion around the double critical point of f_0, i.e., around $z_c = 1$ gives

$$\begin{aligned} f_0(z) &= \frac{1}{2}(z-1)^2 + \mathcal{O}((z-1)^3), \\ f_1(z) &= \xi(z-1) + \mathcal{O}((z-1)^2), \\ f_2(z) &= 0 + \mathcal{O}(z-1). \end{aligned} \tag{90}$$

Fix $r > 1 - \mu_-/\sqrt{T}$ and deform $\Gamma_{0,v}$ to the contour $\gamma = \gamma_1 \cup \gamma_2$ with

$$\gamma_1 = 1 - \mu_-/\sqrt{T} + \mathrm{i}[-r, r], \quad \gamma_2 = \{|z| = r\} \cap \{\operatorname{Re} z < 1 - \mu_-/\sqrt{T}\}. \tag{91}$$

Let us verify that γ is a steep descent path for f_0.

On γ_1 we have $\operatorname{Re} f_0(x + \mathrm{i}y) = x - 1 - \frac{1}{2}\ln(x^2 + y^2)$, so that we have

$$\frac{\mathrm{d} \operatorname{Re} f_0(x+\mathrm{i}y)}{\mathrm{d}y} = -\frac{y}{x^2+y^2}, \qquad y \in [-r, r], \tag{92}$$

with $x = 1 - \mu_-/\sqrt{T}$. Thus f_0 is strictly increasing on $1 - \frac{\mu_-}{\sqrt{T}} + \mathrm{i}[-r, 0)$ and strictly decreasing on $1 - \frac{\mu_-}{\sqrt{T}} + \mathrm{i}(0, r]$. On γ_2 we compute

$$\operatorname{Re} f_0(r\mathrm{e}^{\mathrm{i}\varphi}) = r\cos\varphi - 1 - \ln r, \qquad \varphi \in (\arccos\tfrac{1}{r}, 2\pi - \arccos\tfrac{1}{r}), \tag{93}$$

which means that f_0 is strictly decreasing on $\gamma_2 \cap \{\operatorname{Im} z > 0\}$ and strictly increasing on $\gamma_2 \cap \{\operatorname{Im} z < 0\}$. Thus γ is a steepest descent path for f_0 and the major contribution comes from a line segment $\gamma_\delta = 1 - \frac{\mu_-}{\sqrt{T}} + \mathrm{i}[-\delta,\delta]$ for any $\delta \in (0,1)$. Indeed, the error we make when we integrate along γ_δ instead of γ is of order $\mathcal{O}(\mathrm{e}^{-cT})$ with $c \sim \delta^2$. We therefore consider the integral on γ_δ only,

$$\frac{\sqrt{T}}{2\pi\mathrm{i}} \int_{\gamma_\delta} \mathrm{d}z\, g(z) \mathrm{e}^{\xi\sqrt{T}(z-1) + \frac{\tau T}{2}(z-1)^2} \mathrm{e}^{\mathcal{O}\left(z-1, \sqrt{T}(z-1)^2, T(z-1)^3\right)}. \tag{94}$$

Using $|\mathrm{e}^x - 1| \leq |x|\mathrm{e}^{|x|}$, the difference between (94) and the same integral without the error term can be bounded by

$$\begin{aligned} \frac{\sqrt{T}}{2\pi} \int_{\gamma_\delta} \mathrm{d}z \Big| &\mathrm{e}^{c_1 \xi\sqrt{T}(z-1) + c_2\frac{\tau T}{2}(z-1)^2} \\ &\times \mathcal{O}\left(z-1, \sqrt{T}(z-1)^2, T(z-1)^3, T^{(n-k)/2}(z-1)^{n-k}\right) \Big| \end{aligned} \tag{95}$$

for some constants c_1 and c_2 that can be chosen arbitrarily close to 1 as $\delta \to 0$. By a change of variable $Z = \sqrt{T}(1-z)$ one then sees that this error is of order $\mathcal{O}(T^{-1/2})$. Hence we can consider the integral in (94) without the error term, where the argument of the exponential function simplifies to

$$\frac{\sqrt{T}}{2\pi\mathrm{i}} \int_{\gamma_\delta} \mathrm{d}z\, \mathrm{e}^{\tau T(z-1)^2/2 + \xi\sqrt{T}(z-1)} g(z). \tag{96}$$

The error we make if we extend γ_δ to $1 - \frac{\mu_-}{\sqrt{T}} + \mathrm{i}\mathbb{R}$ is of order $\mathcal{O}(\mathrm{e}^{-cT})$. All together the integral from (88) agrees, up to an error $\mathcal{O}(\mathrm{e}^{-cT}, T^{-1/2})$ uniform in $\xi \in [-L, L]$, with

$$\frac{\sqrt{T}}{2\pi\mathrm{i}} \int_{1 - \frac{\mu_-}{\sqrt{T}} + \mathrm{i}\mathbb{R}} \mathrm{d}z\, \mathrm{e}^{\tau T(z-1)^2/2 + \xi\sqrt{T}(z-1)} g(z), \tag{97}$$

where the poles of g lie on the left of the integration axis. After a change of variable $Z = -\sqrt{T}(z-1)$ this integral can be identified as $\Psi^{n,\tau}_{n-k}(\xi)$.

Finally we show that $\Phi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi') \to \Phi^{n,\tau}_{n-k}(\xi')$. We have

$$\begin{aligned} \Phi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi') = \frac{\sqrt{T}}{2\pi\mathrm{i}} \oint_{\Gamma_v} &\mathrm{d}w\, \mathrm{e}^{\tau T(\ln w - w + 1) - \sqrt{T}\xi' \ln w + n \ln w} \\ &\times \frac{(\sqrt{T}(w-1)+\mu_1)\cdots(\sqrt{T}(w-1)+\mu_{k-1})}{(\sqrt{T}(w-1)+\mu_1)\cdots(\sqrt{T}(w-1)+\mu_n)}, \end{aligned} \tag{98}$$

and by a change of variable $W=-\sqrt{T}(w-1)$ and a Taylor expansion in the exponent we get

$$\begin{aligned}&\Phi^{n,\tau}_{n-k,T,\mathrm{resc}}(\xi')\\&\quad=\frac{(-1)^{n-k+1}}{2\pi\mathrm{i}}\oint_{\Gamma_a}\mathrm{d}W\,\mathrm{e}^{-\tau W^2/2+\xi' W+\mathcal{O}(T^{-1/2})}\,\frac{(w-\mu_1)\cdots(w-\mu_{k-1})}{(w-\mu_1)\cdots(w-\mu_n)},\end{aligned}\tag{99}$$

which converges uniformly for $\xi'\in[-L,L]$ to $\Phi^{n,\tau}_{n-k}(\xi')$. ☐

With the above results we can now prove Theorem 3.

Proof of Theorem 3. Set $m=N(N+1)/2$ and define $n_1,\dots,n_m$ by

$$n_1=1,\;\; n_2=n_3=2,\;\; n_4=n_5=n_6=3,\;\; \dots,\;\; n_{m-N+1}=\dots=n_m=N.\tag{100}$$

For $A\subseteq\mathbb{R}^m$ we set $A_T=(\tau T-\sqrt{T}A)\cap\mathbb{Z}$. Then, we have

$$\begin{aligned}\nu_T(A)&=\sum_{(x_1,\dots,x_m)\in A_T}\det\big[\widetilde{K}^{\mu_T}_{\tau T}((x_i,n_i),(x_j,n_j))\big]_{1\le i,j\le m}\\&=T^{m/2}\int_A\mathrm{d}^m x\,\det\big[\widetilde{K}^{\mu_T}_{\tau T}((\tau T-[x_i]\sqrt{T},n_i),(\tau T-[x_j]\sqrt{T},n_j))\big]_{1\le i,j\le m}\\&=\int_A\mathrm{d}^m x\,\det\big[\widetilde{K}_{\tau,T,\mathrm{resc}}(([x_i],n_i),([x_j],n_j))\big]_{1\le i,j\le m}\end{aligned}\tag{101}$$

and

$$\nu(A)=\int_A\mathrm{d}^m x\,\det\big[K^\mu_\tau((x_i,n_i),(x_j,n_j))\big]_{1\le i,j\le m}.\tag{102}$$

Since the determinants are continuous functions of the kernels, we have by Proposition 4.2 that

$$\begin{aligned}\lim_{T\to\infty}&\det\big[\widetilde{K}^{\mu}_{\tau,T,\mathrm{resc}}(([x_i],n_i),([x_j],n_j))\big]_{1\le i,j\le m}\\&=\det\big[K^\mu_\tau((x_i,n_i),(x_j,n_j))\big]_{1\le i,j\le m}\end{aligned}\tag{103}$$

for all $x_1,\dots,x_m\in\mathbb{R}$. Thus we have shown that the densities of the probability measures in question converge pointwise to each other. Then, (11) is a direct consequence of Scheffé's theorem, see e.g. [3]. ☐

A Determinantal correlations

Since we refer several times to this Lemma 3.4 of [6], we report it here.

Lemma A.1 (Lemma 3.4 of [6]). *Assume we have a signed measure on* $\{x_i^n, n=1,\dots,N, i=1,\dots,n\}$ *given in the form,*

$$\frac{1}{Z_N}\prod_{n=1}^{N-1}\det[\phi_n(x_i^n,x_j^{n+1})]_{1\le i,j\le n+1}\det[\Psi_{N-i}^N(x_j^N)]_{1\le i,j\le N}, \tag{104}$$

where x_{n+1}^n *are some "virtual" variables and* Z_N *is a normalization constant. If* $Z_N\neq 0$*, then the correlation functions are determinantal.*

To write down the kernel we need to introduce some notations. Define

$$\phi^{(n_1,n_2)}(x,y)=\begin{cases}(\phi_{n_1}*\cdots*\phi_{n_2-1})(x,y), & n_1<n_2,\\ 0, & n_1\ge n_2,\end{cases}. \tag{105}$$

where $(a*b)(x,y)=\sum_{z\in\mathbb{Z}}a(x,z)b(z,y)$*, and, for* $1\le n<N$*,*

$$\Psi_{n-j}^n(x):=(\phi^{(n,N)}*\Psi_{N-j}^N)(y),\quad j=1,\dots,N. \tag{106}$$

Set $\phi_0(x_1^0,x)=1$*. Then the functions*

$$\{(\phi_0*\phi^{(1,n)})(x_1^0,x),\dots,(\phi_{n-2}*\phi^{(n-1,n)})(x_{n-1}^{n-2},x),\phi_{n-1}(x_n^{n-1},x)\} \tag{107}$$

are linearly independent and generate the n*-dimensional space* V_n*. Define a set of functions* $\{\Phi_j^n(x), j=0,\dots,n-1\}$ *spanning* V_n *defined by the orthogonality relations*

$$\sum_x\Phi_i^n(x)\Psi_j^n(x)=\delta_{i,j} \tag{108}$$

for $0\le i,j\le n-1$*.*

Under Assumption (A): $\phi_n(x_{n+1}^n,x)=c_n\Phi_0^{(n+1)}(x)$*, for some* $c_n\neq 0$*,* $n=1,\dots,N-1$*, the kernel takes the simple form*

$$K(n_1,x_1;n_2,x_2)=-\phi^{(n_1,n_2)}(x_1,x_2)+\sum_{k=1}^{n_2}\Psi_{n_1-k}^{n_1}(x_1)\Phi_{n_2-k}^{n_2}(x_2). \tag{109}$$

References

[1] M. Adler, E. Nordenstam, and P. van Moerbeke, *Consecutive Minors for Dyson's Brownian Motions*, arXiv:1007.0220 (2010).

[2] Y. Baryshnikov, *GUEs and queues*, Probab. Theory Relat. Fields **119** (2001), 256–274.

[3] P. Billingsley, *Convergence of Probability Measures*, Wiley ed., New York, 1968.

[4] A. Borodin and P. L. Ferrari, *Anisotropic Growth of Random Surfaces in* $2+1$ *Dimensions*, arXiv:0804.3035 (2008).

[5] A. Borodin and P. L. Ferrari, *Large time asymptotics of growth models on space-like paths I: PushASEP*, Electron. J. Probab. **13** (2008), 1380–1418.

[6] A. Borodin, P. L. Ferrari, M. Prähofer, and T. Sasamoto, *Fluctuation Properties of the TASEP with Periodic Initial Configuration*, J. Stat. Phys. **129** (2007), 1055–1080.

[7] A. Borodin and V. Gorin, *Shuffling algorithm for boxed plane partitions*, Adv. Math. **220** (2009), 1739–1770.

[8] A. Borodin and J. Kuan, *Asymptotics of Plancherel measures for the infinite-dimensional unitary group*, Adv. Math. **219** (2008), 894–931.

[9] A. Borodin and J. Kuan, *Random Surface Growth with a Wall and Plancherel Measures for* $O(\infty)$, Comm. Pure Appl. Math. **63** (2010), 831–894.

[10] A. Borodin and S. Péché, *Airy Kernel with Two Sets of Parameters in Directed Percolation and Random Matrix Theory*, J. Stat. Phys. **132** (2008), 275–290.

[11] N. Elkies, G. Kuperbert, M. Larsen, and J. Propp, *Alternating-Sign Matrices and Domino Tilings I and II*, J. Algebr. Comb. **1** (1992), 111–132.

[12] P. L. Ferrari and R. Frings, *On the Partial Connection Between Random Matrices and Interacting Particle Systems*, J. Stat. Phys. **141** (2010), 613–637.

[13] P. J. Forrester and T. Nagao, *Determinantal Correlations for Classical Projection Processes*, J. Stat. Mech. (2011), P08011.

[14] P. J. Forrester and E. Nordenstam, *The Anti-Symmetric GUE Minor Process*, Mosc. Math. J. **9** (2009), 749–774.

[15] V. Gorin and M. Shkolnikov, *Limits of multilevel TASEP and similar processes*, arXiv:1206.3817 (2012).

[16] J. M. Harrison and R. J. Williams, *Multidimensional Reflected Brownian Motions Having Exponential Stationary Distributions*, Ann. Probab. (1987), 115–137.

[17] T. Ichiba and I. Karatzas, *On collisions of Brownian particles*, Ann. Appl. Prob. (2010), 951–977.

[18] K. Johansson and E. Nordenstam, *Eigenvalues of GUE minors*, Electron. J. Probab. **11** (2006), 1342–1371.

[19] I. Karatzas, S. Pal, and M. Shkolnikov, *Systems of Brownian particles with asymmetric collisions*, arXiv:1210.0259 (2012).

[20] A. Metcalfe, *Universality properties of Gelfand-Tsetlin patterns*, Probab. Theory Relat. Fields (2011), online first.

[21] E. Nordenstam, *On the Shuffling Algorithm for Domino Tilings*, Electron. J. Probab. **15** (2010), 75–95.

[22] E. Nordenstam and B. Young, *Domino Shuffling on Novak Half-Hexagons and Aztec Half-Diamonds*, Electr. J. Comb. **18** (2011).

[23] A. Okounkov and N. Reshetikhin, *The Birth of a Random Matrix*, Mosc. Math. J. **6** (2006), 553–566.

[24] S. Pal and J. Pitman, *One-dimensional Brownian particle systems with rank-dependent drifts*, Ann. Appl. Prob. (2008), 2179–2207.

[25] S. R. S. Varadhan and R. J. Williams, *Brownian Motion in a Wedge with Oblique Reflection*, Comm. Pure Appl. Math. (1984), 405–443.

[26] J. Warren, *Dyson's Brownian motions, intertwining and interlacing*, Electron. J. Probab. **12** (2007), 573–590.

[27] J. Warren and P. Windridge, *Some Examples of Dynamics for Gelfand Tsetlin Patterns*, Electron. J. Probab. **14** (2009), 1745–1769.